

purasomes
IRC100+

DERMOAROMA



PROTOCOL FOR
MEN

PROTOCOL FOR INTIMATE REJUVENATION SYSTEM (IRC) FOR MEN

OBJECTIVE

To enhance intimate wellness, improve erectile function, and rejuvenate the male genital area through non-invasive and minimally invasive techniques.

INDICATIONS

- Men seeking improvement in erectile function and penile sensitivity.
- Enhancement of penile skin texture and appearance.
- General rejuvenation and improved blood flow to the genital area.
- Not suitable for patients with visible skin infections, open wounds, or untreated herpes simplex virus.

PRECAUTIONS

- Do not perform the procedure on broken skin, open wounds, or visible infections.
- For patients with a strong history of herpes simplex virus, consider pre-treatment with a prophylactic course of Acyclovir.

CONTRAINDICATIONS

- Active infections in the treatment area.
- History of allergic reactions to any components used in the procedure.
- Severe skin conditions affecting the genital area.

EXPECTED OUTCOME

- Improved penile sensitivity and blood flow.
- Enhanced skin texture and appearance.
- Increased confidence and intimate wellness.

DISCLAIMER:

This protocol should be performed only by qualified medical professionals trained in intimate rejuvenation procedures. Individual results may vary, and all patients should undergo a comprehensive consultation to determine their suitability for the treatment.

PROCEDURE

Step 1: Preparation



- **Apply Numbing Cream:** Use LMX or plain lignocaine cream and apply to the treatment area. Leave on for up to 20 minutes to ensure adequate numbing.
- **Clean the Penis:** Gently cleanse the penis using wet gauze and a non-alcoholic cleaning agent. Allow the area to air dry completely before proceeding.



Step 2: IRC Application

- Apply a thin layer of IRC solution evenly over the penis.
- Ensure complete coverage without excessive pooling of the solution.



Step 3: Dermaroller Application

- Use a dermaroller with a depth of up to 0.5 mm.
- Glide the dermaroller over the penis with moderate and consistent pressure, covering the entire treatment area.

Step 4: Second Layer of IRC

- After dermaroller use, apply another thin layer of IRC over the treated area.



Step 5: Post-Treatment Cleanup

- Gently dab off any excess IRC solution using a clean, dry gauze.
- Allow the treated area to air dry completely before dressing as normal.



Post-Treatment Care

- **Sexual Activity:** Avoid sexual intercourse for up to 48 hours to reduce the risk of infection.
- **Post-Treatment Gel:** Apply the provided post-treatment gel externally to the penis twice daily to promote healing and reduce irritation.

 purasomes
IRC100+



Dermoaroma Srl
Via dei Pirenei 1, Rome RM 00144, Italy
info@dermoaroma.com
dermoaroma.com